

Assessment of chemical properties of tropical peatland soil in Malaysia oil palm plantation

Adesiji Adeolu Richard^{a, d}, Thamer Ahmed Mohammad^a, Nik Norsyahariati Nik Daud^a, Sa'ari Mustapha^b, Alexander Kiew Sayok^c and Gbadebo Anthonia Olukemi^d.

^aDepartment of Civil Engineering, Faculty of Engineering, Universiti Putra, Malaysia, 43400, UPM Serdang, Malaysia. ^bDepartment of Environmental and Chemical Engineering, Faculty of Engineering, Universiti Putra, Malaysia, 43400, UPM Serdang, Malaysia. ^cDepartment of Hydrology and Soil Science, Institute of Biodiversity and Environmental Conservation, Universiti Malaysia, Sarawak, ^d Department of Civil Engineering, Faculty of Engineering, Federal University of Technology, PMB 65, Minna, Nigeria

Abstract

The chemical assessments of the peatland in oil palm plantation in South Selangor Peatland Swamp in Malaysia were evaluated in this study. Soil samples were obtained from fifteen (15) different locations within the study area at three different depths of 0.5m, 1.5 m, and 2.5 m in three replicates at each depth, using peat auger between March and July, 2013 during the secondary maximum rainfall. Parameters evaluated in soil analysis included - pH, moisture content, carbon, nitrogen, sulphur and heavy metals such as manganese, zinc, iron, copper, and phosphorus. Heavy metals were determined using the double acid extraction method while carbon, nitrogen, and sulphur was determined using Trumac CNS Analyzer. From the pH values, the soil close to the surface is more acidic with a mean pH of 3.36 and standard error of 0.15. The mean values of the moisture contents were 363.54 % with SEM of 27.01 and 154.56 % with SEM of 54.64 at 0.5 m and 2.5 m depths respectively. Carbon, had the highest value of 44.27 % at 0.5 m, nitrogen, 0.36 %, sulphur, 0.15 % and heavy metals like manganese, iron, and copper, except zinc, and phosphorus had their mean values either increasing or decreasing with soil depth. Soil carbon was observed to decrease with depth unlike nitrogen and soil pH. All the parameters were observed to either decrease or increase with depth which shows their spatial distribution across the soil mass. This assessment of the tropical peatland soils chemistry has helped in appraising the important roles played by intact peatlands in overall global environmental sustainability.

Keywords: Chemical Properties, Tropical Peatland, Oil Palms Cultivation, Soil Quality, Greenhouse Gases.

E-mail: adrichard01@yahoo.co.uk; ade.richard@futminna.edu.ng

Received: 2015/03/19

Accepted: 2015/11/25

DOI: <http://dx.doi.org/10.4314/njtr.v10i2.3>